

AD-A243 264

2


SECURITY CLASSIFICATION OF THIS PAGE

DTIC
REPORT DOCUMENTForm Approved
OMB No. 0704-0188

1a. REPORT SECURITY CLASSIFICATION Unclassified		1b. RESTRICTIVE MARKINGS	
2a. SECURITY CLASSIFICATION AUTHORITY DEC 7 1991		3. DISTRIBUTION/AVAILABILITY OF REPORT Approved for public release Distribution list enclosed	
2b. DECLASSIFICATION/DOWNGRADING SCHEDULE		5. MONITORING ORGANIZATION REPORT NUMBER(S)	
4. PERFORMING ORGANIZATION REPORT NUMBER(S) Technical Report No. 31		7a. NAME OF MONITORING ORGANIZATION Office of Naval Research	
6a. NAME OF PERFORMING ORGANIZATION State University of New York at Buffalo	6b. OFFICE SYMBOL (If applicable)	7b. ADDRESS (City, State, and ZIP Code) Department of the Navy 800 North Quincy Street Arlington, VA 22217	
6c. ADDRESS (City, State, and ZIP Code) Department of Chemistry Buffalo, NY 14214	9. PROCUREMENT INSTRUMENT IDENTIFICATION NUMBER N00014-90-J-1530; R&T Code 4135002		
8a. NAME OF FUNDING/SPONSORING ORGANIZATION Office of Naval Research	8b. OFFICE SYMBOL (If applicable)	10. SOURCE OF FUNDING NUMBERS	
8c. ADDRESS (City, State, and ZIP Code) Department of the Navy Arlington, VA 22217	PROGRAM ELEMENT NO.	PROJECT NO.	TASK NO. WORK UNIT ACCESSION NO 4135002
11. TITLE (Include Security Classification) Ligand Redistribution Reactions of $\text{Me}_2\text{Ga}(\text{C}_5\text{H}_5)$ and $\text{MeGa}(\text{C}_5\text{H}_5)_2$			
12. PERSONAL AUTHOR(S) O. T. Beachley, Jr., T. L. Royster, Jr. and J. R. Arhar			
13a. TYPE OF REPORT Technical Report	13b. TIME COVERED FROM _____ TO _____	14. DATE OF REPORT (Year, Month, Day) 2 December 1991	15. PAGE COUNT 14
16. SUPPLEMENTARY NOTATION To be published in Journal of Organometallic Chemistry			
17. COSATI CODES		18. SUBJECT TERMS (Continue on reverse if necessary and identify by block number)	
FIELD	GROUP	SUB-GROUP	
		Gallium chemistry, organogallium compounds, cyclopentadienylgallium compounds	
19. ABSTRACT (Continue on reverse if necessary and identify by block number) The compounds $\text{Me}_2\text{Ga}(\text{C}_5\text{H}_5)$ and $\text{MeGa}(\text{C}_5\text{H}_5)_2$ have been prepared by ligand redistribution reactions between appropriate stoichiometric quantities of $\text{Ga}(\text{C}_5\text{H}_5)_3$ and GaMe_3 . Both compounds have been demonstrated by ^1H NMR spectral studies to be unstable in solution and to form symmetrized products by ligand redistribution reactions. Thus, $\text{Me}_2\text{Ga}(\text{C}_5\text{H}_5)$ forms GaMe_3 and $\text{MeGa}(\text{C}_5\text{H}_5)_2$ as primary products whereas $\text{MeGa}(\text{C}_5\text{H}_5)_2$ decomposes to $\text{Ga}(\text{C}_5\text{H}_5)_3$ and $\text{Me}_2\text{Ga}(\text{C}_5\text{H}_5)$. The compound $\text{Me}_2\text{Ga}(\text{C}_5\text{H}_5)$ has also been shown to serve as a cyclopentadienyl transfer reagent as it reacts with FeCl_2 to form $\text{Fe}(\text{C}_5\text{H}_5)_2$ and Me_2GaCl .			
20. DISTRIBUTION/AVAILABILITY OF ABSTRACT <input checked="" type="checkbox"/> UNCLASSIFIED/UNLIMITED <input type="checkbox"/> SAME AS RPT. <input type="checkbox"/> DTIC USERS		21. ABSTRACT SECURITY CLASSIFICATION Unclassified	
22a. NAME OF RESPONSIBLE INDIVIDUAL O. T. Beachley, Jr.		22b. TELEPHONE (Include Area Code) 716-831-3266	22c. OFFICE SYMBOL

OFFICE OF NAVAL RESEARCH
Contract N-00014-90-J-1530
R & T Code 4135002
TECHNICAL REPORT NO. 31

Ligand Redistribution Reactions of



by

O. T. Beachley, Jr., T. L. Royster, Jr. and J. R. Arhar

Prepared for Publication

in

Journal of Organometallic Chemistry

State University of New York at Buffalo
Department of Chemistry
Buffalo, New York 14214

2 December 1991

Reproduction in whole or in part is permitted for
any purpose of the United States Government

*This document has been approved for public release
and sale; its distribution is unlimited

Accession For	
NTIS GRA&I	<input checked="" type="checkbox"/>
DTIC TAB	<input type="checkbox"/>
Unannounced	<input type="checkbox"/>
Justification	
By	
Distribution/	
Availability Codes	
Dist	Avail and/or Special
A-1	




91

91-17174



Contribution from the Department of Chemistry
State University of New York at Buffalo, Buffalo, NY 14214

Ligand Redistribution Reactions of



by

O. T. Beachley, Jr., T. L. Royster, Jr. and J. R. Arhar

Summary:

The compounds $\text{Me}_2\text{Ga}(\text{C}_5\text{H}_5)$ and $\text{MeGa}(\text{C}_5\text{H}_5)_2$ have been prepared by ligand redistribution reactions between appropriate stoichiometric quantities of $\text{Ga}(\text{C}_5\text{H}_5)_3$ and GaMe_3 . Both compounds have been demonstrated by ^1H NMR spectral studies to be unstable in solution and to form symmetrized products by ligand redistribution reactions. Thus, $\text{Me}_2\text{Ga}(\text{C}_5\text{H}_5)$ forms GaMe_3 and $\text{MeGa}(\text{C}_5\text{H}_5)_2$ as primary products whereas $\text{MeGa}(\text{C}_5\text{H}_5)_2$ decomposes to $\text{Ga}(\text{C}_5\text{H}_5)_3$ and $\text{Me}_2\text{Ga}(\text{C}_5\text{H}_5)$. The compound $\text{Me}_2\text{Ga}(\text{C}_5\text{H}_5)$ has also been shown to serve as a cyclopentadienyl transfer reagent as it reacts with FeCl_2 to form $\text{Fe}(\text{C}_5\text{H}_5)_2$ and Me_2GaCl .

Introduction

Heteroleptic organometallic compounds of the heavier group 13 elements gallium and indium form an interesting class of compounds which have not been extensively investigated. The first example of a fully characterized organoindium compound¹ with two different alkyl substituents $[\text{In}(\text{CH}_2\text{CMe}_3)(\text{CH}_2\text{SiMe}_3)\text{Cl}]_2$ was only recently reported. Even though this indium compound could be prepared, isolated and characterized, the related organogallium compounds¹ $\text{Ga}(\text{Me})(\text{CH}_2\text{CMe}_3)\text{Cl}$ and $\text{Ga}(\text{CH}_2\text{CMe}_3)(\text{CH}_2\text{CMe}_2\text{Ph})\text{Cl}$ could not be isolated. All data were consistent with the isolation of symmetrized products, GaR_2Cl and $\text{GaR}'_2\text{Cl}$. Even though the synthesis of pure samples of $\text{Ga}(\text{Me})(\text{CH}_2\text{CMe}_3)\text{Cl}$ and $\text{Ga}(\text{CH}_2\text{CMe}_3)(\text{CH}_2\text{CMePh})\text{Cl}$ were unsuccessful, other heteroleptic organogallium compounds including $\text{Me}_2\text{Ga}(\text{C}_5\text{H}_5)$,^{2,3,4} $\text{Et}_2\text{Ga}(\text{C}_5\text{H}_5)$,^{2,3} $\text{Me}_2\text{Ga}(\text{C CPh})$,^{5,6} $\text{Me}_2\text{Ga}(\text{C}\equiv\text{CMe})$,⁷ $\text{Me}_2\text{Ga}(\text{CH}=\text{CH}_2)$,⁸ $\text{Me}_2\text{Ga}(\text{t-Bu})$ ⁹ and $\text{MeGa}(\text{t-Bu})_2$ ⁹ have been reported. However, since we were unable to prepare pure samples of unsymmetrically substituted organogallium compounds, we were prompted to investigate the detailed nature of $\text{Me}_2\text{Ga}(\text{C}_5\text{H}_5)$ in the solution phase.

The compound $\text{Me}_2\text{Ga}(\text{C}_5\text{H}_5)$ ² has been reported as a colorless, crystalline solid with a melting point of 68-69 °C. The compound had been prepared from Me_2GaCl with $\text{Na}(\text{K})\text{C}_5\text{H}_5$ in cyclohexane at 50-70 °C and had been isolated by sublimation at 65 °C (0.5 mm Hg).^{2,3} The elemental analysis for carbon was observed to be significantly lower than the calculated value whereas the percent gallium was higher than the calculated value.¹ The compound was also reported to have an association number of 1.3-1.4 in benzene solution.² The X-ray structural study⁴ defined chains of dimethylgallium groups bridged by cyclopentadienyl rings at the 1 and 3 positions. The availability of $\text{Ga}(\text{C}_5\text{H}_5)_3$ ¹⁰ and GaMe_3 in our laboratories as well as the facility of ligand

distribution reactions for preparing pure heteroleptic compounds prompted us to attempt the preparation and isolation of $\text{Me}_2\text{Ga}(\text{C}_5\text{H}_5)$ and $\text{MeGa}(\text{C}_5\text{H}_5)_2$. The preparation of $\text{Me}_2\text{Ga}(\text{C}_5\text{H}_5)$ as a pure compound with excellent analyses for carbon and hydrogen was achieved. However, the melting point and association number in benzene solution were significantly different than those previously reported. In addition, NMR spectral studies of solutions of isolated samples $\text{Me}_2\text{Ga}(\text{C}_5\text{H}_5)$, $\text{MeGa}(\text{C}_5\text{H}_5)_2$ and of solutions of these compounds prepared directly from GaMe_3 and $\text{Ga}(\text{C}_5\text{H}_5)_3$ suggest the occurrence of ligand redistribution reactions. Thus, available data suggest that neither $\text{Me}_2\text{Ga}(\text{C}_5\text{H}_5)$ nor $\text{MeGa}(\text{C}_5\text{H}_5)_2$ exist as single compounds in solution.

Experimental

All compounds described in this investigation were extremely sensitive to oxygen and moisture and were manipulated in a standard vacuum line or under a purified argon atmosphere. The starting material $\text{Ga}(\text{C}_5\text{H}_5)_3$ was prepared and purified by the literature method¹⁰ whereas GaMe_3 was purchased from Strem Chemicals and used as received. Solvents were dried by conventional procedures. Elemental analyses were performed by Schwarzkopf Microanalytical Laboratory, Woodside, NY. The ^1H NMR spectra were recorded either at 90 MHz by using a Varian EM390 spectrometer or at 400 MHz by using a Varian VXR-400 spectrometer. Proton chemical shifts are reported in δ units (ppm) and are referenced to SiMe_4 at δ 0.00 ppm and C_6H_6 at δ 7.15 ppm. All samples for NMR spectra were contained in sealed NMR tubes. Melting points were observed in sealed capillaries. Molecular weights were measured cryoscopically in benzene by using an instrument similar to that described by Shriver and Drezdson.¹¹

Synthesis of $\text{Me}_2\text{Ga}(\text{C}_5\text{H}_5)$. A 0.689 g (2.60 mmol) sample of $\text{Ga}(\text{C}_5\text{H}_5)_3$ was dissolved in 50 mL of pentane. Then, 0.597 g (5.20 mmol) of GaMe_3 was condensed into the reaction flask at -196°C . As the reaction mixture was warmed to room temperature, a colorless precipitate formed. After 2 h at room temperature, the solvent was removed by vacuum distillation. The reaction vessel was connected to a 100 mL Schlenk flask by using a glass elbow joint. The crude product was then sublimed at 60°C into the Schlenk flask that was cooled to -196°C to yield 1.20 g (7.28 mmol, 93.4%) of $\text{Me}_2\text{Ga}(\text{C}_5\text{H}_5)$ as a colorless crystalline solid. $\text{Me}_2\text{Ga}(\text{C}_5\text{H}_5)$: mp $97-98^\circ\text{C}$. ^1H NMR (CCl_4 , 90 MHz) δ 6.29 (s, C_5H_5 , 5 H), -0.31 (s, Me, 6 H). (C_6H_6 , 90 MHz) δ 6.22 (s, C_5H_5 , 5 H), -0.42 (s, Me, 6 H); (C_6D_6 , 400 MHz) 6.19 (s, C_5H_5), -0.46 (s), -1.13 (s,br) (see Results and Discussion); (d^8 -THF, 90 MHz)

δ 5.95 (s, C_5H_5), -0.54 (s, Me, $GaMe_3 \cdot THF$), -0.68 (s, Me, $Me_2Ga(C_5H_5) \cdot THF$), -1.11 (s, Me, $MeGa(C_5H_5)_2 \cdot THF$). IR (Nujol mull, cm^{-1}) 3100 (w), 3080 (w), 1776 (w), 1720 (w), 1620 (w), 1399 (m), 1360 (m), 1342 (m), 1192 (m), 1181 (m), 1068 (w), 1056 (w), 1002 (w), 984 (m), 879 (w), 872 (m), 854 (m), 836 (s), 813 (m), 797 (s), 779 (s), 745 (s), 717 (s), 602 (m), 578 (vs), 522 (m), 290 (w). Cryoscopic molecular weight, benzene solution, formula weight 164.9 (obsd. molality, obsd. mol. wt., association): 0.0811, 178, 1.08; 0.0734, 181, 1.10; 0.0730, 177, 1.08; 0.0588, 179, 1.08; 0.0496, 178, 1.08. Anal. Calcd: C, 50.99; H, 6.72. Found: C, 50.82; H, 6.76.

Reaction of 2 Mol of $Ga(C_5H_5)_3$ per Mol of $GaMe_3$. A 0.132 g (1.15 mmol) sample of $GaMe_3$ was vacuum distilled into a reaction flask containing 0.610 g (2.30 mmol) of $Ga(C_5H_5)_3$ in 30 mL of pentane. The reaction mixture was stirred at room temperature for 16 h. Then the solvent was removed by vacuum distillation. The yellow liquid which remained in the reaction flask was identified by 1H NMR spectroscopy as $MeGa(C_5H_5)_2$. After standing at room temperature for 16 h, colorless crystals were observed in the liquid. The compounds $MeGa(C_5H_5)_2$ and $Me_2Ga(C_5H_5)$ were identified by the 1H NMR spectrum of the resulting liquid-solid mixture dissolved in CCl_4 . Yellow Liquid. 1H NMR (CCl_4 , 90 MHz) δ 6.25 (s, C_5H_5 , 10H, $MeGa(C_5H_5)_2$), -1.04 (s, Me, 3 H, $MeGa(C_5H_5)_2$). Liquid-Solid Mixture. 1H NMR (CCl_4 , 90 MHz) δ 6.20 (s, C_5H_5), -0.33 (s, Me, $Me_2Ga(C_5H_5)$), -1.12 (s, Me, $MeGa(C_5H_5)_2$).

Synthesis of $MeGa(C_5H_5)_2 \cdot THF$. A 0.111 g (0.967 mmol) sample of $GaMe_3$ was vacuum distilled into a reaction flask containing 0.513 g (1.93 mmol) of $Ga(C_5H_5)_3$ in 25 mL of pentane. The reaction mixture was stirred at room temperature for 16 h. Then, 10 mL of THF was vacuum distilled into the flask and the reaction mixture was stirred for an additional 3 h at room temperature. After the solvent was removed in vacuo, a slightly yellow

liquid remained in the flask. ^1H NMR (d^6 -benzene, 90 MHz) δ 6.20 (s, C_5H_5), 3.29 (m, THF), 1.18 (m, THF), -0.33 (s, Me), -0.91 (s, Me). (THF, 90 MHz) δ 5.94 (s, C_5H_5), -0.64 (s, Me, $\text{Me}_2\text{Ga}(\text{C}_5\text{H}_5)\cdot\text{THF}$), -1.13 (s, Me, $\text{MeGa}(\text{C}_5\text{H}_5)_2\cdot\text{THF}$). Anal. Calcd for $\text{MeGa}(\text{C}_5\text{H}_5)_2\cdot\text{THF}$: C, 62.76; H, 7.37. Found: C, 62.56; H, 7.17.

^1H NMR Spectrum of a Mixture of $\text{Me}_2\text{Ga}(\text{C}_5\text{H}_5)$ and GaMe_3 . A sample of 0.0317 g (0.194 mmol) of $\text{Me}_2\text{Ga}(\text{C}_5\text{H}_5)$ was placed in an NMR tube. Then, a small quantity of GaMe_3 (~ 0.07 mmol) and C_6D_6 as solvent were added by vacuum distillation. The ^1H NMR spectrum at 400 MHz of the resulting solution revealed lines at -0.31 (singlet) and -6.20 ppm (singlet). (See Results and Discussion).

^1H NMR Spectrum of Equimolar Mixtures of $\text{Ga}(\text{C}_5\text{H}_5)_3$ and GaMe_3 in Different Solvents. The nature of the exchange reaction in the different solvents, CCl_4 , C_6H_6 and THF, was investigated by ^1H NMR spectroscopy by combining $\text{Ga}(\text{C}_5\text{H}_5)_3$ and GaMe_3 in a 1:1 mol ratio in the given solvent. In a typical experiment, a weighed quantity of $\text{Ga}(\text{C}_5\text{H}_5)_3$ was placed in a reaction vessel connected to an NMR tube. Then GaMe_3 was condensed into the reaction vessel at -196 °C. The ^1H NMR spectrum (90 MHz) of the reaction mixture was recorded immediately upon warming the solution to room temperature.

CCl_4 . The reaction mixture was prepared from 0.200 g (0.755 mmol) of $\text{Ga}(\text{C}_5\text{H}_5)_3$ and 0.0867 g (0.755 mmol) of GaMe_3 in 3 mL of CCl_4 . ^1H NMR: -1.06 (very broad singlet, Me, $\text{MeGa}(\text{C}_5\text{H}_5)_2$, relative intensity 1), -0.30 (very broad singlet, Me, $\text{Me}_2\text{Ga}(\text{C}_5\text{H}_5)$, relative intensity 2), 6.20 (singlet, C_5H_5).

C_6H_6 . The reaction mixture was prepared from 0.0612 g (0.231 mmol) sample of $\text{Ga}(\text{C}_5\text{H}_5)_3$ and 0.0265 g (0.231 mmol) of GaMe_3 in 3 mL of C_6H_6 . ^1H NMR: -1.07 (broadened singlet, Me, $\text{MeGa}(\text{C}_5\text{H}_5)_2$, relative intensity 1),

-0.42 (broadened singlet, Me, $\text{Me}_2\text{Ga}(\text{C}_5\text{H}_5)$, relative intensity 2), 6.20 (singlet, C_5H_5).

THF. The reaction mixture was prepared from 0.0550 g (0.207 mmol) of $\text{Ga}(\text{C}_5\text{H}_5)_3$ and 0.0236 g (0.207 mmol) of GaMe_3 in 3 mL of THF. ^1H NMR: -1.21 (sharp singlet, Me, $\text{MeGa}(\text{C}_5\text{H}_5)_2\cdot\text{THF}$, relative intensity 1), -0.68 (sharp singlet, Me, $\text{Me}_2\text{Ga}(\text{C}_5\text{H}_5)\cdot\text{THF}$, relative intensity 2), 5.95 (singlet, C_5H_5).

Synthesis of $\text{Fe}(\text{C}_5\text{H}_5)_2$: Procedure A: A reaction mixture of 0.250 g (1.51 mmol) of $\text{Me}_2\text{Ga}(\text{C}_5\text{H}_5)$ and 0.0960 g (0.726 mmol) of FeCl_2 in 25 mL of benzene was stirred at room temperature for 16 h. After the solvent was removed in vacuo, an orange solid remained in the reaction flask. Then, a volatile colorless crystalline product was sublimed at room temperature into a -196°C trap. This colorless solid was identified as Me_2GaCl by its ^1H NMR spectrum in benzene solution (δ 0.23, Me; δ 0.23 Me_2GaCl). The remaining orange product was sublimed at 45°C under high vacuum. The solid was washed several times with dilute aqueous HCl and distilled water to remove any trace of Me_2GaCl . After drying the product in the air, 0.0638 g of $\text{Fe}(\text{C}_5\text{H}_5)_2$ (0.343 mmol, 47.2% based on FeCl_2) was isolated.

Procedure B: In the drybox, a 0.500 g (3.03 mmol) sample of $\text{Me}_2\text{Ga}(\text{C}_5\text{H}_5)$ was placed into a 100 mL Schlenk flask and 0.200 g (1.58 mmol) of FeCl_2 was contained in a connecting side-arm dumper. Then, 50 mL of THF was vacuum distilled onto $\text{Me}_2\text{Ga}(\text{C}_5\text{H}_5)$ and 25 mL of the solvent was distilled onto the FeCl_2 . The sparingly soluble FeCl_2 was added with stirring to the solution of $\text{Me}_2\text{Ga}(\text{C}_5\text{H}_5)$ at room temperature. After 3 h the solid FeCl_2 had disappeared and an orange solution was observed. The reaction mixture was stirred for a total of 16 h at room temperature. The solvent was removed in vacuo and an orange solid mixed with a liquid remained in the reaction flask. By heating the liquid-solid mixture with a 40°C oil bath, the

product was transferred under high vacuum into a Schlenk flask cooled to $-196\text{ }^{\circ}\text{C}$. With the product mixture maintained at $-196\text{ }^{\circ}\text{C}$, distilled water was added to the flask. After slowly warming the reaction mixture to room temperature, dilute aqueous HCl was added. The solution was then filtered and the orange solid product was collected on a glass frit. The product $\text{Me}_2\text{GaOH}\cdot\text{THF}$ was identified by its ^1H NMR spectrum ($\delta -0.32$). The orange solid was washed several more times with distilled water to remove any trace of $\text{Me}_2\text{GaOH}\cdot\text{THF}$. After drying the product in the air, 0.210 g (1.13 mmol, 74.6% based on $\text{Me}_2\text{Ga}(\text{C}_5\text{H}_5)$) of $\text{Fe}(\text{C}_5\text{H}_5)_2$ was isolated.


Results and Discussion

The compound $\text{Me}_2\text{Ga}(\text{C}_5\text{H}_5)$ has been prepared in nearly quantitative yield by a ligand redistribution reaction between GaMe_3 and $\text{Ga}(\text{C}_5\text{H}_5)_3$ in a 2:1 mol ratio in pentane. It is noteworthy that GaMe_3 and $\text{Ga}(\text{C}_5\text{H}_5)_3$ are soluble in pentane but $\text{Me}_2\text{Ga}(\text{C}_5\text{H}_5)$ precipitates from the solution as it is formed. The product, as demonstrated by its elemental analyses for carbon and hydrogen and its melting point, is significantly more pure than the product originally isolated from the metathesis reaction between Me_2GaCl and $\text{Na}(\text{K})\text{C}_5\text{H}_5$ in cyclohexane.² The observed carbon and hydrogen analyses for the new sample differed from the calculated values by only -0.17% and +0.04%, respectively. The melting point of the compound prepared by the ligand redistribution reaction, 97-98 °C, was considerably higher than that reported for the sample prepared by the metathetical reaction,² 68-69 °C. These data suggest that the compound as originally reported was impure. A possible impurity might be unreacted Me_2GaCl . The Me_2GaCl might not be consumed completely if the $\text{Na}(\text{K})\text{Cl}$ formed in the metathesis reaction coated the insoluble reagent $\text{Na}(\text{K})\text{C}_5\text{H}_5$ and thereby hindered reaction. It is also possible that the insoluble nature of $\text{Me}_2\text{Ga}(\text{C}_5\text{H}_5)$ in aliphatic hydrocarbon solvents might hinder the preparative reaction. Since $\text{Me}_2\text{Ga}(\text{C}_5\text{H}_5)$ and Me_2GaCl sublime at similar temperatures, unreacted Me_2GaCl would not have been removed and a product with a low melting point and low analytical data for percent carbon as well as high percent gallium would result.

The cryoscopic molecular weight studies of pure $\text{Me}_2\text{Ga}(\text{C}_5\text{H}_5)$ dissolved in benzene provide results which are in disagreement with the earlier work.² The product from the ligand redistribution reaction had an average association value of 1.1 which was independent of concentration in the range of 0.08 to 0.05 m. These data suggest the presence of monomeric species in

benzene solution. The original impure product had a reported association value of 1.3 - 1.4 (no concentration range given).² An impurity of Me_2GaCl which exists as a dimer in benzene solution would produce an observed molecular weight which would be higher than calculated. Thus, the degree of association of $\text{Me}_2\text{Ga}(\text{C}_5\text{H}_5)$ is similar to most other triorganogallium compounds including $\text{Ga}(\text{C}_5\text{H}_5)_3$ ¹⁰ which are monomeric in benzene solution. Only $\text{Me}_2\text{Ga}(\text{C}\equiv\text{CPh})$ ⁵ and $\text{Me}_2\text{Ga}(\text{C}\equiv\text{CMe})$ ⁷ have been reported as dimers.


Proton NMR (400 MHz) studies of benzene solutions of $\text{Me}_2\text{Ga}(\text{C}_5\text{H}_5)$ suggest the occurrence of a ligand redistribution reaction to form symmetrized products (Equation 1) and of methyl and cyclopentadienyl group



exchange reactions between reactants and products. The spectrum of $\text{Me}_2\text{Ga}(\text{C}_5\text{H}_5)$ exhibits three lines, a sharp line at 6.19 ppm, a slightly broadened line at -0.46 ppm and a significantly broader but low intensity line centered at -1.13 ppm. (The line at -1.13 ppm was not observed in 90 MHz spectra). The line at 6.19 ppm can be assigned to cyclopentadienyl group protons. The line at -0.46 ppm arises from methyl group protons on $\text{Me}_2\text{Ga}(\text{C}_5\text{H}_5)$ and GaMe_3 , which are averaged by methyl group exchange reactions. A separate line at -0.19 ppm, as expected for GaMe_3 , was not observed. The line at -1.13 ppm is probably related to the protons of the methyl group of $\text{MeGa}(\text{C}_5\text{H}_5)_2$ which, in turn, are averaged by exchange with GaMe_3 formed by the ligand redistribution reaction (Equation 1). The suggestion of a methyl group exchange reaction between $\text{Me}_2\text{Ga}(\text{C}_5\text{H}_5)$ and GaMe_3 was verified by an independent experiment. When $\text{Me}_2\text{Ga}(\text{C}_5\text{H}_5)$ and GaMe_3 were mixed in benzene, only one methyl-gallium and one cyclopentadienyl group

resonance was observed. The broad line at -1.13 ppm originally observed in the spectrum of $\text{Me}_2\text{Ga}(\text{C}_5\text{H}_5)$ and assigned to the methyl protons of $\text{MeGa}(\text{C}_5\text{H}_5)_2$ disappeared. In addition, the methyl-gallium line moved from -0.46 to -0.31 ppm as would be expected if methyl group exchange were occurring in C_6D_6 . Thus, $\text{Me}_2\text{Ga}(\text{C}_5\text{H}_5)$ does not exist as a single compound in benzene solution. Ligand redistribution reactions to form symmetrized products occur in solution. The observation that pure $\text{Me}_2\text{Ga}(\text{C}_5\text{H}_5)$ can be isolated from the preparative reaction in pentane must be related to the precipitation of the compound from solution as it is formed. The bridge bonds between cyclopentadienyl groups to form four coordinate gallium⁴ must be important to the stability of the compound as a solid.


Attempts have been made to prepare, isolate, and characterize the first example of a heavier group 13 compound with two cyclopentadienyl groups, $\text{MeGa}(\text{C}_5\text{H}_5)_2$. When the reagents $\text{Ga}(\text{C}_5\text{H}_5)_3$ and GaMe_3 were combined in a 2:1 ratio in pentane, a clear solution formed. When the pentane was removed, a yellow liquid was obtained. This yellow liquid was dissolved in turn in CCl_4 and it exhibited a ^1H NMR spectrum (100 MHz) which was appropriate for a compound with the formula $\text{MeGa}(\text{C}_5\text{H}_5)_2$, i.e., one line for C_5H_5 at 6.25 ppm and one line for Me at -1.04 ppm with an integration ratio of 10:3. However, in the absence of solvent, the yellow liquid deposited a crystalline solid after 16 h. The ^1H NMR spectrum of this liquid-solid mixture dissolved in CCl_4 exhibited three lines, i.e., one C_5H_5 line at 6.20 ppm and two Me lines at -0.33 and -1.12 ppm. The chemical shift of the line at -0.33 ppm is indicative of $\text{Me}_2\text{Ga}(\text{C}_5\text{H}_5)$ (^1H NMR lines at 6.29 and -0.31 ppm). These data are consistent with the occurrence of a symmetrization reaction (Equation 2). Thus, $\text{MeGa}(\text{C}_5\text{H}_5)_2$ decomposed slowly and could not be fully characterized.



The ligand redistribution reactions of $\text{Me}_2\text{Ga}(\text{C}_5\text{H}_5)$ and $\text{MeGa}(\text{C}_5\text{H}_5)_2$ were confirmed by ^1H NMR spectra of solutions formed upon addition of the Lewis base, THF. The ^1H NMR spectrum of $\text{Me}_2\text{Ga}(\text{C}_5\text{H}_5)$ in THF exhibited four lines at -1.11, -0.68, -0.54 and 5.95 ppm. The three lines at higher field are assigned to the methyl group protons in $\text{MeGa}(\text{C}_5\text{H}_5)_2 \cdot \text{THF}$, $\text{Me}_2\text{Ga}(\text{C}_5\text{H}_5) \cdot \text{THF}$ and $\text{Me}_3\text{Ga} \cdot \text{THF}$, respectively. The equilibrium constant for Equation 3 calculated




from the integration values of the lines was 0.13. The ^1H NMR spectrum of a sample of $\text{MeGa}(\text{C}_5\text{H}_5)_2$ prepared in pentane as previously described and then dissolved in THF had three lines. These lines are consistent with the




assignments given above and with the redistribution equilibrium as shown in Equation 4. The equilibrium constant calculated from NMR integration data for Equation 4 was 0.17. The compound $\text{Me}_2\text{Al}(\text{C}_5\text{H}_5) \cdot \text{THF}$ ¹² is reported to undergo an analogous redistribution equilibrium in THF. However, the product $\text{MeAl}(\text{C}_5\text{H}_5)_2 \cdot \text{THF}$ has never been isolated and studied.

The ligand redistribution reaction between $\text{Ga}(\text{C}_5\text{H}_5)_3$ and GaMe_3 was investigated further by combining the reagents in a 1:1 ratio in THF, C_6H_6 and CCl_4 and by studying the ^1H NMR spectra of the resulting solutions. All data are consistent with the occurrence of ligand redistribution reactions



to form $\text{MeGa}(\text{C}_5\text{H}_5)$ and $\text{MeGa}(\text{C}_5\text{H}_5)_2$ (Equation 5). Only three ^1H NMR lines, as expected, were observed in each spectrum. However, it is noteworthy that the rates of exchange for the reactions in each solvent were different. The lines for the THF solution were very sharp and were indicative of slow exchange. In contrast, two extremely broad methyl-gallium lines were observed for the CCl_4 solution. The lines observed for benzene solution had widths which were in between those observed for THF and CCl_4 solutions, but certainly not sharp. The chemical shifts of the two methyl-gallium lines assigned to $\text{Me}_2\text{Ga}(\text{C}_5\text{H}_5)$ and $\text{MeGa}(\text{C}_5\text{H}_5)_2$ are consistent with all previous discussions.

The compound $\text{Me}_2\text{Ga}(\text{C}_5\text{H}_5)$ has been utilized as a source of C_5H_5 units for the synthesis of ferrocene. The ferrocene was characterized by its



melting point and ^1H NMR spectrum whereas the colorless crystalline product was identified as Me_2GaCl by ^1H NMR spectroscopy. All characterization data for the two products were consistent with data for authentic samples.

Acknowledgements. This work was supported in part by the Office of Naval Research and by a generous grant from Eastman Kodak Co. The authors also thank Sun-Hua Chao for preparing solutions and recording the ^1H NMR spectra at 400 MHz.

References

1. O. T. Beachey, Jr., J. D. Maloney, M. R. Churchill, C. H. Lake, *Organometallics* 10, (1991), page .
2. J. Stadelhofer, J. Weidlein, A. Haaland, *J. Organomet. Chem.*, 84, (1975), C1.
3. J. Stadelhofer, J. Weidlein, P. Fischer, A. Haaland, *J. Organomet. Chem.*, 116, (1976), 55.
4. K. Mertz, F. Zettler, H. D. Hausen, J. Weidlein, *J. Organomet. Chem.*, 122, (1976), 159.
5. E. A. Jeffrey, T. Mole, *J. Organomet. Chem.*, 11, (1968), 393.
6. B. Tecle, W. H. Ilsley, J. P. Oliver, *Inorg. Chem.* 20, (1981), 2335.
7. W. Fries, W. Schwarz, H.-D. Hausen, J. Weidlein, *J. Organomet. Chem.*, 159, (1978), 373.
8. W. Fries, K. Sille, J. Weidlein, A. Haaland, *Spectrochim. Acta*, 36A, (1980), 611.
9. W. M. Cleaver, A. R. Barron, *Chemtronics*, 4, (1989), 146.
10. O. T. Beachley, Jr.; T. D. Getman, R. U. Kirss, R. B. Hallock, W. E. Hunter, J. L. Atwood, *Organometallics* 4, (1985), 751.
11. D. F. Shriver, M. A. Drezdson, "The Manipulation of Air Sensitive Compounds", Wiley: New York, (1986), p. 38.
12. W. R. Kroll, *J. Chem. Soc. Chem. Commun.*, (1969), 844.

TECHNICAL REPORT DISTRIBUTION LIST - GENERAL

Office of Naval Research (2)
Chemistry Division, Code 1113
800 North Quincy Street
Arlington, Virginia 22217-5000

Dr. Richard W. Drisko (1)
Naval Civil Engineering
Laboratory
Code L52
Port Hueneme, CA 93043

Dr. James S. Murday (1)
Chemistry Division, Code 6100
Naval Research Laboratory
Washington, D.C. 20375-5000

Dr. Harold H. Singerman (1)
David Taylor Research Center
Code 283
Annapolis, MD 21402-5067

Dr. Robert Green, Director (1)
Chemistry Division, Code 385
Naval Weapons Center
China Lake, CA 93555-6001

Chief of Naval Research (1)
Special Assistant for Marine
Corps Matters
Code 00MC
800 North Quincy Street
Arlington, VA 22217-5000

Dr. Eugene C. Fischer (1)
Code 2840
David Taylor Research Center
Annapolis, MD 21402-5067

Defense Technical Information
Center (2)
Building 5, Cameron Station
Alexandria, VA 22314

Dr. Elek Lindner (1)
Naval Ocean Systems Center
Code 52
San Diego, CA 92152-5000

Commanding Officer (1)
Naval Weapons Support Center
Dr. Bernard E. Doua
Crane, Indiana 47522-5050

* Number of copies to forward